A NOTEWORTHY COLLECTION OF ANDIRA GALEOTTIANA

BY

RICHARD EVANS SCHULTES

Among other plants recently sent from the Botanical Museum to Dr. Paul C. Standley for study, there was a specimen of the rare *Andira Galeottiana* belonging to the Leguminosae.

Andira Galeottiana Standley in Contrib. U.S. Nat. Herb. 20 (1919) 217.

Mexico: Oaxaca, District of Choapam, San Juan Lalana, long. 95° 45′, lat. 17°25′, alt. 600 m., May 10, 1939, Schultes & Reko 852 (Econ. Herb. Oakes Ames No. 3930; Herb. Gray; Herb. Field Mus. Nat. Hist.).

The type of Andira Galeottiana, an endemic of the District of Choapam (Oaxaca) and adjacent parts of Vera Cruz, was collected in Catemaco, Vera Cruz (Nelson 424). A second collection was cited in the type description as "collected at Lalana (Chinantla) Puebla, July, 1844, by H. Galeotti (no. 3464)". This citation should be corrected to read: "Lalana (Chinantla), Oaxaca..." Andira Galeottiana has not been collected in Puebla so far as herbarium records indicate.

Schultes & Reko 852 was collected in San Juan Lalana, the same village where Galeotti, nearly a century ago, found the cotype. It is apparently the third collection of this tree.

In Lalana, Andira Galeottiana is called by the Chinantee name mo-dzaw and by the Zapotee name gia-bia. In 1844, Galeotti recorded the Spanish vernacular name as macayo, and in 1939, Reko and I found the same name still in use for this plant. The species is abundant in Lalana where its profusion of rose-purple panicles is very conspicuous on the hillsides in May and June.

[200]